45 IN-2

CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Safety glasses, gloves and protective clothing should be worn when using this product.

DISCLAIMER/CONDITIONS OF SALE: **1)** Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. **2)** The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. **3)** The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. **4)** The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. **5)** Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. **6)** Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

BATCH No.

9 333056 001891

SPRAY GRO®

LIQUID FERTILIZER

CHARGE

N P K

13 : 14 : 9 w/v

20 Litre NET ☐ 200 Litre NET ☐ 1000 Litre NET ☐

BROADACRE: Foliar rate of application per min water/hectare: 50 litres.

CROP	Fertigation L/Ha	Foliar L/Ha	
CEREALS	30-50	N/A	Apply through soil injection at planting.
COTTON	25-30	N/A	Apply through soil injection at planting. Reduce rates 5-6L/ha) with Anhydrous ammonia, if has been injected. Fertigate again at early flowering.
CANOLA	20-30	N/A	Apply at planting in conjunction with 42N.
LEGUMES	20-25	N/A	Apply at planting.
MAIZE/ SUGARCANE	20-30	N/A	Apply at planting in maize and sugarcane. Apply every ratoon crop.
LUCERNE	N/A	3-5	Apply when sufficient leaf area exists after every cut.

HORTICULTURE: Foliar rate of application per min water/hectare: 150-350 litres. Tree crops may require higher rates (500-1000L).

CROP	Fertigation L/Ha	Foliar L/Ha	
AVOCADOES	7-10	2-3	Apply to juvenile trees at early establishment-fertigation. Apply foliar at least two weeks before flowering.
BANANAS	8-12	N/A	Apply early in the season. Repeat if necessary.
BEANS	4-6	N/A	Apply at 2 leaf stage. Repeat if necessary.
BRASSICAS	6-10	2-3	Apply through soil after transplant. Apply foliar at early heading.
CUCURBITS	5-10	2-3	Apply at planting. Foliar at fruit set. Repeat if necessary.
LEAFY GREENS	7-10	N/A	Apply every 14 days from mid-crop through to harvest.
OLIVE TREES	15-18	3-5	Apply early season through soil. Foliar from early through to late fruiting. Fertigate after harvest.
PAW PAW	8-10	1-3	Apply through soil early in the season. Foliar application 21 days after flowering as required.
PEANUTS	20-25	N/A	Apply at planting.
PINEAPPLES	6-10	1-3	Apply early season. Foliar 14 days before fruit set. Repeat if necessary.
POTATOES	5-10	2-4	Apply at planting - foliar.
RED WINE GRAPES	10-20	3-5	Fertigate after bud burst. Foliar, apply before flowering. Repeat at early fruiting.
STRAWBERRIES	8-12	N/A	Apply at planting. Repeat 14-21 day interval as required.
WHITE WINE GRAPES	8-15	2-3	Fertigate after bud break. Foliar when sufficient leaf area develops. Repeat before flowering. Fertigage post harvest.
TOMATOES	7-10	N/A	Apply after transplant. Repeat before truss formation, if required.
TABLE GRAPES	10-15	N/A	Apply at bud break. Repeat at fruiting.
TURF	N/A	300-500ml/ 100m²	Apply at regular intervals on established turf.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/W %
Nitrogen N	13	9.55
Phosphorus P	14	P_2O_5 23.57
Potassium K	9	K_2O 7.94

Specific gravity 1.36

STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

drumMUSTER eligible container

NOT APPLICABLE TO 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA